

Sign Variations and Descents

Aram Dermenjian

Joint with: Nantel Bergeron and John Machacek



York University

11 January 2021

Sign vectors

- A *sign vector* is a vector whose components are signs.
- \mathcal{V}_n is the set of all sign vectors.
- Think of these as vectors associated to coordinate in \mathbb{R}^n where we only look at signs.



Example



Sign vectors

- A *sign vector* is a vector whose components are signs.
- \mathcal{V}_n is the set of all sign vectors.
- Think of these as vectors associated to coordinate in \mathbb{R}^n where we only look at signs.




Example



Sign vectors

- A *sign vector* is a vector whose components are signs.
- \mathcal{V}_n is the set of all sign vectors.
- Think of these as vectors associated to coordinate in \mathbb{R}^n where we only look at signs.


Example



Sign vectors

- A *sign vector* is a vector whose components are signs.
- \mathcal{V}_n is the set of all sign vectors.
- Think of these as vectors associated to coordinate in \mathbb{R}^n where we only look at signs.



Example



Sign vectors

- A *sign vector* is a vector whose components are signs.
- \mathcal{V}_n is the set of all sign vectors.
- Think of these as vectors associated to coordinate in \mathbb{R}^n where we only look at signs.


Example



Sign vectors


- A *sign vector* is a vector whose components are signs.
- \mathcal{V}_n is the set of all sign vectors.
- Think of these as vectors associated to coordinate in \mathbb{R}^n where we only look at signs.

Example




Real projective space

- *Real Projective space \mathbb{RP}^n* is \mathbb{R}^{n+1} without the origin and where we associate each vector with all scalar multiples.



Real projective space - \mathbb{RP}^1



Images from: math.stackexchange.com by Zev Chonoles

Projective sign vectors

- Let \mathcal{PV}_n be the set of sign vectors in \mathbb{RP}^{n-1} .
- In other words, for $\omega \in \mathcal{V}_n$, then $\omega \sim \omega'$ if and only if $\omega = \omega'$ or $\omega = -\omega'$.

Example

$$\begin{aligned}\mathcal{V}_1 &= \{(+), (0), (-)\} & \mathcal{V}_2 &= \{(+, +), (+, 0), (+, -), \\ &&& (0, +), (0, 0), (0, -), \\ &&& (-, +), (-, 0), (-, -)\}\end{aligned}$$

Projective sign vectors

- Let \mathcal{PV}_n be the set of sign vectors in \mathbb{RP}^{n-1} .
- In other words, for $\omega \in \mathcal{V}_n$, then $\omega \sim \omega'$ if and only if $\omega = \omega'$ or $\omega = -\omega'$.

Example

$$\begin{array}{ll}\mathcal{V}_1 = \{(+), (0), (-)\} & \mathcal{V}_2 = \{(+, +), (+, 0), (+, -), \\ & \quad \downarrow \\ & \quad (0, +), (0, 0), (0, -), \\ \{(+), (0), (-)\} & \quad (-, +), (-, 0), (-, -)\}\end{array}$$

Projective sign vectors

- Let \mathcal{PV}_n be the set of sign vectors in \mathbb{RP}^{n-1} .
- In other words, for $\omega \in \mathcal{V}_n$, then $\omega \sim \omega'$ if and only if $\omega = \omega'$ or $\omega = -\omega'$.

Example

$$\begin{array}{ll}\mathcal{V}_1 = \{(+), (0), (-)\} & \mathcal{V}_2 = \{(+, +), (+, 0), (+, -), \\ & \quad \downarrow \\ & \quad (0, +), (0, 0), (0, -), \\ \{(+), (\textcolor{red}{0}), (-)\} & \quad (-, +), (-, 0), (-, -)\}\end{array}$$

Projective sign vectors

- Let \mathcal{PV}_n be the set of sign vectors in \mathbb{RP}^{n-1} .
- In other words, for $\omega \in \mathcal{V}_n$, then $\omega \sim \omega'$ if and only if $\omega = \omega'$ or $\omega = -\omega'$.

Example

$$\begin{array}{ll}\mathcal{V}_1 = \{(+), (0), (-)\} & \mathcal{V}_2 = \{(+, +), (+, 0), (+, -), \\ & \quad (0, +), (0, 0), (0, -), \\ & \quad (-, +), (-, 0), (-, -)\} \\ \downarrow & \\ \{\textcolor{blue}{(+)}, \textcolor{red}{(-)}\} & \end{array}$$

Projective sign vectors

- Let \mathcal{PV}_n be the set of sign vectors in \mathbb{RP}^{n-1} .
- In other words, for $\omega \in \mathcal{V}_n$, then $\omega \sim \omega'$ if and only if $\omega = \omega'$ or $\omega = -\omega'$.

Example

$$\begin{array}{ll}\mathcal{V}_1 = \{(+), (0), (-)\} & \mathcal{V}_2 = \{(+, +), (+, 0), (+, -), \\ & \qquad \qquad \qquad \downarrow \\ & \qquad \qquad \qquad (0, +), (0, 0), (0, -), \\ \mathcal{PV}_1 = \{(+)\} & \qquad \qquad \qquad (-, +), (-, 0), (-, -)\}\end{array}$$

Projective sign vectors

- Let \mathcal{PV}_n be the set of sign vectors in \mathbb{RP}^{n-1} .
- In other words, for $\omega \in \mathcal{V}_n$, then $\omega \sim \omega'$ if and only if $\omega = \omega'$ or $\omega = -\omega'$.

Example

$$\mathcal{V}_1 = \{(+), (0), (-)\} \quad \mathcal{V}_2 = \{(+, +), (+, 0), (+, -),$$

↓

$$\mathcal{PV}_1 = \{(+)\}$$

$$(0, +), (0, 0), (0, -),$$

$$(-, +), (-, 0), (-, -)\}$$

↓

$$\{(+, +), (+, 0), (+, -),$$

$$(0, +), (0, 0), (0, -),$$

$$(-, +), (-, 0), (-, -)\}$$

Projective sign vectors

- Let \mathcal{PV}_n be the set of sign vectors in \mathbb{RP}^{n-1} .
- In other words, for $\omega \in \mathcal{V}_n$, then $\omega \sim \omega'$ if and only if $\omega = \omega'$ or $\omega = -\omega'$.

Example

$$\mathcal{V}_1 = \{(+), (0), (-)\} \quad \mathcal{V}_2 = \{(+, +), (+, 0), (+, -),$$

↓

$$\mathcal{PV}_1 = \{(+)\}$$

$$(0, +), (0, 0), (0, -),$$

$$(-, +), (-, 0), (-, -)\}$$

↓

$$\{(+, +), (+, 0), (+, -),$$

$$(0, +), (\cancel{0}, \cancel{0}), (0, -),$$

$$(-, +), (-, 0), (-, -)\}$$

Projective sign vectors

- Let \mathcal{PV}_n be the set of sign vectors in \mathbb{RP}^{n-1} .
- In other words, for $\omega \in \mathcal{V}_n$, then $\omega \sim \omega'$ if and only if $\omega = \omega'$ or $\omega = -\omega'$.

Example

$$\mathcal{V}_1 = \{(+), (0), (-)\} \quad \mathcal{V}_2 = \{(+, +), (+, 0), (+, -),$$

↓

$$\mathcal{PV}_1 = \{(+)\}$$

$$(0, +), (0, 0), (0, -),$$

$$(-, +), (-, 0), (-, -)\}$$

↓

$$\{(+, +), (+, 0), (+, -),$$

$$(0, +), (0, -),$$

$$(-, +), (-, 0), (\underline{\underline{-}}, \underline{\underline{-}})\}$$

Projective sign vectors

- Let \mathcal{PV}_n be the set of sign vectors in \mathbb{RP}^{n-1} .
- In other words, for $\omega \in \mathcal{V}_n$, then $\omega \sim \omega'$ if and only if $\omega = \omega'$ or $\omega = -\omega'$.

Example

$$\mathcal{V}_1 = \{(+), (0), (-)\} \quad \mathcal{V}_2 = \{(+, +), (+, 0), (+, -),$$

↓

$$\mathcal{PV}_1 = \{(+)\}$$

$$(0, +), (0, 0), (0, -),$$

$$(-, +), (-, 0), (-, -)\}$$

↓

$$\{(+, +), \color{blue}{(+, 0)}, (+, -),$$

$$(0, +), (0, -),$$

$$(-, +), \color{red}{(-, 0)}\}$$

Projective sign vectors

- Let \mathcal{PV}_n be the set of sign vectors in \mathbb{RP}^{n-1} .
- In other words, for $\omega \in \mathcal{V}_n$, then $\omega \sim \omega'$ if and only if $\omega = \omega'$ or $\omega = -\omega'$.

Example

$$\mathcal{V}_1 = \{(+), (0), (-)\} \quad \mathcal{V}_2 = \{(+, +), (+, 0), (+, -),$$

↓

$$\mathcal{PV}_1 = \{(+)\}$$

$$(0, +), (0, 0), (0, -),$$

$$(-, +), (-, 0), (-, -)\}$$

↓

$$\{(+, +), (+, 0), (+, -),$$

$$(0, +), (0, -),$$

$$(\textcolor{red}{-, +})\}$$

Projective sign vectors

- Let \mathcal{PV}_n be the set of sign vectors in \mathbb{RP}^{n-1} .
- In other words, for $\omega \in \mathcal{V}_n$, then $\omega \sim \omega'$ if and only if $\omega = \omega'$ or $\omega = -\omega'$.

Example

$$\begin{array}{ll}\mathcal{V}_1 = \{(+), (0), (-)\} & \mathcal{V}_2 = \{(+, +), (+, 0), (+, -), \\ & \quad \downarrow \\ & \quad (0, +), (0, 0), (0, -), \\ \mathcal{PV}_1 = \{(+)\} & \quad (-, +), (-, 0), (-, -)\} \\ & \quad \downarrow \\ & \quad \{(+, +), (+, 0), (+, -), \\ & \quad \textcolor{blue}{(0, +)}, \textcolor{red}{(0, -)}\}\end{array}$$

Projective sign vectors

- Let \mathcal{PV}_n be the set of sign vectors in \mathbb{RP}^{n-1} .
- In other words, for $\omega \in \mathcal{V}_n$, then $\omega \sim \omega'$ if and only if $\omega = \omega'$ or $\omega = -\omega'$.

Example

$$\mathcal{V}_1 = \{(+), (0), (-)\} \quad \mathcal{V}_2 = \{(+, +), (+, 0), (+, -),$$

\downarrow

$$\mathcal{PV}_1 = \{(+)\} \quad \mathcal{V}_2 = \{(0, +), (0, 0), (0, -),$$

\downarrow

$$\mathcal{PV}_2 = \{(+, +), (+, 0), (+, -), (0, +)\}$$

Projective sign vectors

- Let \mathcal{PV}_n be the set of sign vectors in \mathbb{RP}^{n-1} .
- In other words, for $\omega \in \mathcal{V}_n$, then $\omega \sim \omega'$ if and only if $\omega = \omega'$ or $\omega = -\omega'$.

$\mathcal{PV}_n \cong \{\omega \in \mathcal{V}_n : \text{First non-zero entry of } \omega \text{ is } +\}$.

Example

$$\mathcal{V}_1 = \{(+), (0), (-)\}$$



$$\mathcal{PV}_1 = \{(+)\}$$

$$\mathcal{V}_2 = \{(+, +), (+, 0), (+, -),$$

$$(0, +), (0, 0), (0, -),$$

$$(-, +), (-, 0), (-, -)\}$$



$$\mathcal{PV}_2 = \{(+, +), (+, 0), (+, -), (0, +)\}$$


Partially ordered set

Given a set P and a relation \preccurlyeq , then (P, \preccurlyeq) is a *partially ordered set (poset)* if for all $a, b, c \in P$:

- $a \preccurlyeq a$ (reflexive)
- $a \preccurlyeq b$ and $b \preccurlyeq a$ implies $a = b$ (anti-symmetric)
- $a \preccurlyeq b$ and $b \preccurlyeq c$ implies $a \preccurlyeq c$ (transitive)

Example

The poset $(\mathbb{N}, |)$:



Ordering projective sign vectors

- Let P_n denote the poset $(\mathcal{PV}_n, <)$ where for $\omega, \omega' \in \mathcal{PV}_n$:

$$\omega' < \omega \iff \pm\omega' \subseteq \omega$$

in other words, if either ω' or $-\omega'$ is obtained from ω by replacing some components with 0.

Example

$$(+, +) \quad (+, -)$$

$$(+, 0) \quad (0, +)$$

Ordering projective sign vectors

- Let P_n denote the poset $(\mathcal{PV}_n, <)$ where for $\omega, \omega' \in \mathcal{PV}_n$:

$$\omega' < \omega \iff \pm\omega' \subseteq \omega$$

in other words, if either ω' or $-\omega'$ is obtained from ω by replacing some components with 0.

Example

$$(+, +) \quad (+, -)$$

$$(+, 0) \quad (0, +)$$

Ordering projective sign vectors

- Let P_n denote the poset $(\mathcal{PV}_n, <)$ where for $\omega, \omega' \in \mathcal{PV}_n$:

$$\omega' < \omega \iff \pm\omega' \subseteq \omega$$

in other words, if either ω' or $-\omega'$ is obtained from ω by replacing some components with 0.

Example

$$(+, +) \qquad (+, -)$$



$$(+, 0) \qquad (0, +)$$

Ordering projective sign vectors

- Let P_n denote the poset $(\mathcal{PV}_n, <)$ where for $\omega, \omega' \in \mathcal{PV}_n$:

$$\omega' < \omega \iff \pm\omega' \subseteq \omega$$

in other words, if either ω' or $-\omega'$ is obtained from ω by replacing some components with 0.

Example

$$(+, +) \qquad (+, -)$$



$$(+, 0) \qquad (0, +)$$


Ordering projective sign vectors

- Let P_n denote the poset $(\mathcal{PV}_n, <)$ where for $\omega, \omega' \in \mathcal{PV}_n$:

$$\omega' < \omega \iff \pm\omega' \subseteq \omega$$

in other words, if either ω' or $-\omega'$ is obtained from ω by replacing some components with 0.

Example




Ordering projective sign vectors

- Let P_n denote the poset $(\mathcal{PV}_n, <)$ where for $\omega, \omega' \in \mathcal{PV}_n$:

$$\omega' < \omega \iff \pm\omega' \subseteq \omega$$

in other words, if either ω' or $-\omega'$ is obtained from ω by replacing some components with 0.

Example




Ordering projective sign vectors

- Let P_n denote the poset $(\mathcal{PV}_n, <)$ where for $\omega, \omega' \in \mathcal{PV}_n$:

$$\omega' < \omega \iff \pm\omega' \subseteq \omega$$

in other words, if either ω' or $-\omega'$ is obtained from ω by replacing some components with 0.

Example




Ordering projective sign vectors

- Let P_n denote the poset $(\mathcal{PV}_n, <)$ where for $\omega, \omega' \in \mathcal{PV}_n$:

$$\omega' < \omega \iff \pm\omega' \subseteq \omega$$

in other words, if either ω' or $-\omega'$ is obtained from ω by replacing some components with 0.

Example




Ordering projective sign vectors

- Let P_n denote the poset $(\mathcal{PV}_n, <)$ where for $\omega, \omega' \in \mathcal{PV}_n$:

$$\omega' < \omega \iff \pm\omega' \subseteq \omega$$

in other words, if either ω' or $-\omega'$ is obtained from ω by replacing some components with 0.

Example




Ordering projective sign vectors

- Let P_n denote the poset $(\mathcal{PV}_n, <)$ where for $\omega, \omega' \in \mathcal{PV}_n$:

$$\omega' < \omega \iff \pm\omega' \subseteq \omega$$

in other words, if either ω' or $-\omega'$ is obtained from ω by replacing some components with 0.

Example




Ordering projective sign vectors

- Let P_n denote the poset $(\mathcal{PV}_n, <)$ where for $\omega, \omega' \in \mathcal{PV}_n$:

$$\omega' < \omega \iff \pm\omega' \subseteq \omega$$

in other words, if either ω' or $-\omega'$ is obtained from ω by replacing some components with 0.

Example




Ordering projective sign vectors

- Let P_n denote the poset $(\mathcal{PV}_n, <)$ where for $\omega, \omega' \in \mathcal{PV}_n$:

$$\omega' < \omega \iff \pm\omega' \subseteq \omega$$

in other words, if either ω' or $-\omega'$ is obtained from ω by replacing some components with 0.

Example




Ordering projective sign vectors

- Let P_n denote the poset $(\mathcal{PV}_n, <)$ where for $\omega, \omega' \in \mathcal{PV}_n$:

$$\omega' < \omega \iff \pm\omega' \subseteq \omega$$

in other words, if either ω' or $-\omega'$ is obtained from ω by replacing some components with 0.

Example




Order complex (of a poset)

- *Simplicial complex Δ* - A collection of sets s.t. $\sigma \in \Delta$ and $\tau \subseteq \sigma$ implies $\tau \in \Delta$.
- The sets are called *faces*. Maximal sets are called *facets*.
- *Order complex $\Delta(P)$* of a poset P - Simplicial complex where faces are chains in P .

Example

(+, +) (+, -)




(+, 0) (0, +)

P_2

Order complex (of a poset)

- *Simplicial complex Δ* - A collection of sets s.t. $\sigma \in \Delta$ and $\tau \subseteq \sigma$ implies $\tau \in \Delta$.
- The sets are called *faces*. Maximal sets are called *facets*.
- *Order complex $\Delta(P)$* of a poset P - Simplicial complex where faces are chains in P .


Example



Order complex (of a poset)

- *Simplicial complex Δ* - A collection of sets s.t. $\sigma \in \Delta$ and $\tau \subseteq \sigma$ implies $\tau \in \Delta$.
- The sets are called *faces*. Maximal sets are called *facets*.
- *Order complex $\Delta(P)$* of a poset P - Simplicial complex where faces are chains in P .


Example



Order complex (of a poset)

- *Simplicial complex Δ* - A collection of sets s.t. $\sigma \in \Delta$ and $\tau \subseteq \sigma$ implies $\tau \in \Delta$.
- The sets are called *faces*. Maximal sets are called *facets*.
- *Order complex $\Delta(P)$* of a poset P - Simplicial complex where faces are chains in P .



Example



Order complex (of a poset)

- *Simplicial complex Δ* - A collection of sets s.t. $\sigma \in \Delta$ and $\tau \subseteq \sigma$ implies $\tau \in \Delta$.
- The sets are called *faces*. Maximal sets are called *facets*.
- *Order complex $\Delta(P)$* of a poset P - Simplicial complex where faces are chains in P .

Example




Order complex (of a poset)

- *Simplicial complex Δ* - A collection of sets s.t. $\sigma \in \Delta$ and $\tau \subseteq \sigma$ implies $\tau \in \Delta$.
- The sets are called *faces*. Maximal sets are called *facets*.
- *Order complex $\Delta(P)$* of a poset P - Simplicial complex where faces are chains in P .


Example

(+, +) (+, -)



(+, 0) (0, +)

P_2



Order complex (of a poset)

- *Simplicial complex Δ* - A collection of sets s.t. $\sigma \in \Delta$ and $\tau \subseteq \sigma$ implies $\tau \in \Delta$.
- The sets are called *faces*. Maximal sets are called *facets*.
- *Order complex $\Delta(P)$* of a poset P - Simplicial complex where faces are chains in P .

Example


(+, +) (+, -)

$$\left\{ \begin{array}{l} (+,+) \\ | \\ (0,+) \end{array} \right\}$$

$$\left\{ \begin{array}{l} (+,+) \\ | \\ (+,0) \end{array} \right\}$$

$$\left\{ \begin{array}{l} (+,-) \\ | \\ (+,0) \end{array} \right\}$$

(+, 0) (0, +)



P_2

$$\{(0,+)\}$$

$$\{(+,+)\}$$

$$\{(+,-)\}$$

$$\{(+,0)\}$$


$$\{\emptyset\}$$

Order complex (of a poset)

- *Simplicial complex Δ* - A collection of sets s.t. $\sigma \in \Delta$ and $\tau \subseteq \sigma$ implies $\tau \in \Delta$.
- The sets are called *faces*. Maximal sets are called *facets*.
- *Order complex $\Delta(P)$* of a poset P - Simplicial complex where faces are chains in P .

Example

$(+, +)$ $(+, -)$



$(+, 0)$ $(0, +)$

P_2

$$\left\{ \begin{matrix} (+, +) \\ | \\ (0, +) \end{matrix} \right\} \quad \left\{ \begin{matrix} (+, -) \\ | \\ (0, +) \end{matrix} \right\} \quad \left\{ \begin{matrix} (+, +) \\ | \\ (+, 0) \end{matrix} \right\} \quad \left\{ \begin{matrix} (+, -) \\ | \\ (+, 0) \end{matrix} \right\}$$



$$\{(0, +)\} \quad \{(+, +)\} \quad \{(+, -)\} \quad \{(+, 0)\}$$

$$\{\emptyset\}$$

Order complex (of a poset)

- *Simplicial complex Δ* - A collection of sets s.t. $\sigma \in \Delta$ and $\tau \subseteq \sigma$ implies $\tau \in \Delta$.
- The sets are called *faces*. Maximal sets are called *facets*.
- *Order complex $\Delta(P)$* of a poset P - Simplicial complex where faces are chains in P .


Example



f-vector

- Δ a d -dim simplicial complex.
- f_i = number of i -dim faces
- *f-vector* is vector faces: $f(\Delta) = (f_{-1}, f_0, f_1, \dots, f_d)$.
- $f(\Delta(P))$ is number of elements in each row.

Example



$$f(\Delta(P_2)) = (1, 4, 4)$$


h-vectors

- Δ a d -dim simplicial complex with $f(\Delta) = (f_{-1}, f_0, \dots, f_d)$.

$$h_k = \sum_{i=0}^k (-1)^{k-i} \binom{d-i}{k-i} f_{i-1}.$$

- *h-vector* is vector of h_k s: $h(\Delta) = (h_0, h_1, \dots, h_{d+1})$.

Example



How can we find the h -vector?

Theorem (Stanley 1975)

If a simplicial complex Δ is Cohen-Macaulay, its h -vector has nonnegative entries.

Theorem (Machacek 2019)

The order complex $\Delta(P_n)$ is Cohen-Macaulay.

Questions

- *Is there a nice way to compute the h -vector of $\Delta(P_n)$?*

Partitionable simplicial complex

Conjecture (Stanley 1979, Garsia 1980; Counterexample
Duval, Goeckner, Klivans, Martin 2016)

Every Cohen-Macaulay simplicial complex is partitionable.

Proposition (Stanley)

If Δ is partitionable, then the partitioning gives us the h -vector.


Partitionable

A simplicial complex Δ is *partitionable* if

$$\Delta = \bigsqcup [G_i, F_i]$$

where F_i is a facet.

Example




Partitionable

A simplicial complex Δ is *partitionable* if

$$\Delta = \bigsqcup [G_i, F_i]$$

where F_i is a facet.

Example




Partitionable

A simplicial complex Δ is *partitionable* if

$$\Delta = \bigsqcup [G_i, F_i]$$

where F_i is a facet.

Example




Partitionable

A simplicial complex Δ is *partitionable* if

$$\Delta = \bigsqcup [G_i, F_i]$$

where F_i is a facet.

Example




Partitionable

A simplicial complex Δ is *partitionable* if

$$\Delta = \bigsqcup [G_i, F_i]$$

where F_i is a facet.

Example



Partitionable


Proposition (Stanley)

Let Δ be a partitionable simplicial complex with partitioning $\Delta = \sqcup [G_i, F_i]$ where F_i is a facet. Then

$$h_i(\Delta) = |\{j : |G_j| = i\}|.$$

Example

$$h(\Delta(P_2)) = (h_0, h_1, h_2)$$



$$\Delta(P_2)$$

Partitionable


Proposition (Stanley)

Let Δ be a partitionable simplicial complex with partitioning $\Delta = \sqcup [G_i, F_i]$ where F_i is a facet. Then

$$h_i(\Delta) = |\{j : |G_j| = i\}|.$$

Example

$$h(\Delta(P_2)) = (1, h_1, h_2)$$



$$\Delta(P_2)$$

Partitionable


Proposition (Stanley)

Let Δ be a partitionable simplicial complex with partitioning $\Delta = \sqcup [G_i, F_i]$ where F_i is a facet. Then

$$h_i(\Delta) = |\{j : |G_j| = i\}|.$$

Example

$$h(\Delta(P_2)) = (1, 2, h_2)$$



Partitionable


Proposition (Stanley)

Let Δ be a partitionable simplicial complex with partitioning $\Delta = \sqcup [G_i, F_i]$ where F_i is a facet. Then

$$h_i(\Delta) = |\{j : |G_j| = i\}|.$$

Example

$$h(\Delta(P_2)) = (1, 2, 1)$$



Main Theorem

Let D_n be a type D Coxeter group and let des_B denote the type B descent set of an element $\pi \in D_n$.

Theorem (Bergeron, D., Machacek 2020BP)

The order complex $\Delta(P_n)$ is partitionable. Moreover,

$$h_i(\Delta(P_n)) = |\{\pi \in D_n : |\text{des}_B(\pi)| = i\}|$$

for each $0 \leq i \leq n$.

Coxeter groups

Type A_n

The elements in type A_n Coxeter groups can be represented as permutations in \mathfrak{S}_{n+1} .

$$\mathfrak{S}_{n+1}.$$

$$57238146 \in A_7$$

Type B_n

The elements in type B_n Coxeter groups can be represented as *signed* permutations of \mathfrak{S}_n .

$$5\bar{7}23\bar{8}\bar{1}46 \in B_8$$

Type D_n

The elements in type D_n Coxeter groups can be represented as *even signed* permutations of \mathfrak{S}_n .

$$5\bar{7}2\bar{3}\bar{8}\bar{1}46 \in D_8$$

Descents - Type A

For $\pi = \pi_1 \dots \pi_{n+1} \in A_n$ let $\text{des}_A(\pi)$ denote the descent set of π .

$$\text{des}_A(\pi) = \{i : \pi_i > \pi_{i+1} \text{ for } 1 \leq i \leq n\}$$

Example

$$\pi = 57238146 \in A_7$$

Descents - Type A

For $\pi = \pi_1 \dots \pi_{n+1} \in A_n$ let $\text{des}_A(\pi)$ denote the descent set of π .

$$\text{des}_A(\pi) = \{i : \pi_i > \pi_{i+1} \text{ for } 1 \leq i \leq n\}$$

Example

$$\begin{aligned}\pi = & \color{red}{57}238146 \in A_7 \\ & \color{red}{12}345678\end{aligned}$$

$$\text{des}_A(\pi) = \emptyset$$

Descents - Type A

For $\pi = \pi_1 \dots \pi_{n+1} \in A_n$ let $\text{des}_A(\pi)$ denote the descent set of π .

$$\text{des}_A(\pi) = \{i : \pi_i > \pi_{i+1} \text{ for } 1 \leq i \leq n\}$$

Example

$$\pi = 5 \color{red}{7} \color{black} 2 \color{black} 3 \color{black} 8 \color{black} 1 \color{black} 4 \color{black} 6 \in A_7$$

$$\color{red}{1} \color{black} 2 \color{black} 3 \color{black} 4 \color{black} 5 \color{black} 6 \color{black} 7 \color{black} 8$$

$$\text{des}_A(\pi) = \emptyset$$

Descents - Type A

For $\pi = \pi_1 \dots \pi_{n+1} \in A_n$ let $\text{des}_A(\pi)$ denote the descent set of π .

$$\text{des}_A(\pi) = \{i : \pi_i > \pi_{i+1} \text{ for } 1 \leq i \leq n\}$$

Example

$$\begin{aligned}\pi = & \color{red}{5} \color{black}7 \color{red}{2} \color{black}3 \color{black}8 \color{black}1 \color{black}4 \color{black}6 \in A_7 \\ & \color{red}{1} \color{black}2 \color{red}{3} \color{black}4 \color{black}5 \color{black}6 \color{black}7 \color{black}8\end{aligned}$$

$$\text{des}_A(\pi) = \{2\}$$

Descents - Type A

For $\pi = \pi_1 \dots \pi_{n+1} \in A_n$ let $\text{des}_A(\pi)$ denote the descent set of π .

$$\text{des}_A(\pi) = \{i : \pi_i > \pi_{i+1} \text{ for } 1 \leq i \leq n\}$$

Example

$$\pi = 57\mathbf{23}8146 \in A_7$$

$$12\mathbf{34}5678$$

$$\text{des}_A(\pi) = \{2\}$$

Descents - Type A

For $\pi = \pi_1 \dots \pi_{n+1} \in A_n$ let $\text{des}_A(\pi)$ denote the descent set of π .

$$\text{des}_A(\pi) = \{i : \pi_i > \pi_{i+1} \text{ for } 1 \leq i \leq n\}$$

Example

$$\pi = 572\mathbf{38}146 \in A_7$$

$$123\mathbf{45}678$$

$$\text{des}_A(\pi) = \{2\}$$

Descents - Type A

For $\pi = \pi_1 \dots \pi_{n+1} \in A_n$ let $\text{des}_A(\pi)$ denote the descent set of π .

$$\text{des}_A(\pi) = \{i : \pi_i > \pi_{i+1} \text{ for } 1 \leq i \leq n\}$$

Example

$$\begin{array}{c} \pi = 57238146 \in A_7 \\ 12345678 \end{array}$$

$$\text{des}_A(\pi) = \{2\}$$

Descents - Type A

For $\pi = \pi_1 \dots \pi_{n+1} \in A_n$ let $\text{des}_A(\pi)$ denote the descent set of π .

$$\text{des}_A(\pi) = \{i : \pi_i > \pi_{i+1} \text{ for } 1 \leq i \leq n\}$$

Example

$$\pi = 5723\mathbf{81}46 \in A_7$$

$$1234\mathbf{56}78$$

$$\text{des}_A(\pi) = \{2, 5\}$$

Descents - Type A

For $\pi = \pi_1 \dots \pi_{n+1} \in A_n$ let $\text{des}_A(\pi)$ denote the descent set of π .

$$\text{des}_A(\pi) = \{i : \pi_i > \pi_{i+1} \text{ for } 1 \leq i \leq n\}$$

Example

$$\pi = 57238\mathbf{146} \in A_7$$

$$12345\mathbf{678}$$

$$\text{des}_A(\pi) = \{2, 5\}$$

Descents - Type A

For $\pi = \pi_1 \dots \pi_{n+1} \in A_n$ let $\text{des}_A(\pi)$ denote the descent set of π .

$$\text{des}_A(\pi) = \{i : \pi_i > \pi_{i+1} \text{ for } 1 \leq i \leq n\}$$

Example

$$\pi = 572381\textcolor{red}{4}6 \in A_7$$

$$123465\textcolor{red}{7}8$$

$$\text{des}_A(\pi) = \{2, 5\}$$

Descents - Type A

For $\pi = \pi_1 \dots \pi_{n+1} \in A_n$ let $\text{des}_A(\pi)$ denote the descent set of π .

$$\text{des}_A(\pi) = \{i : \pi_i > \pi_{i+1} \text{ for } 1 \leq i \leq n\}$$

Example

$$\pi = 57238146 \in A_7$$

$$\text{des}_A(\pi) = \{2, 5\}$$

Descents - Type *B*

For $\pi = \pi_1 \dots \pi_n \in B_n$ let $\text{des}_B(\pi)$ denote the descent set of π .

$$\text{des}_B(\pi) = \{i : \pi_i > \pi_{i+1} \text{ for } 0 \leq i < n\}$$

where $\pi_0 = 0$.

Example

$$\pi = 5\bar{7}2\bar{3}\bar{8}\bar{1}46 \in B_8$$

Descents - Type *B*

For $\pi = \pi_1 \dots \pi_n \in B_n$ let $\text{des}_B(\pi)$ denote the descent set of π .

$$\text{des}_B(\pi) = \{i : \pi_i > \pi_{i+1} \text{ for } 0 \leq i < n\}$$

where $\pi_0 = 0$.

Example

$$\pi = 5\bar{7}2\bar{3}\bar{8}\bar{1}46 \in B_8$$

To find descent, we add a 0 in front, and calculate like “normal”.

$$05\bar{7}2\bar{3}\bar{8}\bar{1}46$$

$$012345678$$

$$\text{des}_B(\pi) = \emptyset$$

Descents - Type B

For $\pi = \pi_1 \dots \pi_n \in B_n$ let $\text{des}_B(\pi)$ denote the descent set of π .

$$\text{des}_B(\pi) = \{i : \pi_i > \pi_{i+1} \text{ for } 0 \leq i < n\}$$

where $\pi_0 = 0$.

Example

$$\pi = 5\bar{7}2\bar{3}\bar{8}\bar{1}46 \in B_8$$

$$0\color{red}{5}\bar{7}2\bar{3}\bar{8}\bar{1}46$$

$$0\color{red}{1}2345678$$

$$\text{des}_B(\pi) = \{1\}$$

Descents - Type B

For $\pi = \pi_1 \dots \pi_n \in B_n$ let $\text{des}_B(\pi)$ denote the descent set of π .

$$\text{des}_B(\pi) = \{i : \pi_i > \pi_{i+1} \text{ for } 0 \leq i < n\}$$

where $\pi_0 = 0$.

Example

$$\pi = 5\bar{7}2\bar{3}\bar{8}\bar{1}46 \in B_8$$

$$05\bar{7}2\bar{3}\bar{8}\bar{1}46$$

$$012\textcolor{red}{34}5678$$

$$\text{des}_B(\pi) = \{1, 3\}$$

Descents - Type B

For $\pi = \pi_1 \dots \pi_n \in B_n$ let $\text{des}_B(\pi)$ denote the descent set of π .

$$\text{des}_B(\pi) = \{i : \pi_i > \pi_{i+1} \text{ for } 0 \leq i < n\}$$

where $\pi_0 = 0$.

Example

$$\pi = 5\bar{7}2\bar{3}\bar{8}\bar{1}46 \in B_8$$

$$05\bar{7}2\bar{3}\bar{8}\bar{1}46$$

$$0123\textcolor{red}{45}678$$

$$\text{des}_B(\pi) = \{1, 3, 4\}$$

Descents - Type B

For $\pi = \pi_1 \dots \pi_n \in B_n$ let $\text{des}_B(\pi)$ denote the descent set of π .

$$\text{des}_B(\pi) = \{i : \pi_i > \pi_{i+1} \text{ for } 0 \leq i < n\}$$

where $\pi_0 = 0$.

Example

$$\pi = 5\bar{7}2\bar{3}\bar{8}\bar{1}46 \in B_8$$

$$\text{des}_B(\pi) = \{1, 3, 4\}$$

Descents - Type D

For $\pi = \pi_1 \dots \pi_n \in D_n$ let $\text{des}_D(\pi)$ denote the descent set of π .

$$\text{des}_D(\pi) = \{i : \pi_i > \pi_{i+1} \text{ for } 0 \leq i < n\}$$

where $\pi_0 = -\pi_2$.

Example

$$\pi = 5\bar{7}2\bar{3}\bar{8}\bar{1}46 \in D_8$$

Descents - Type D

For $\pi = \pi_1 \dots \pi_n \in D_n$ let $\text{des}_D(\pi)$ denote the descent set of π .

$$\text{des}_D(\pi) = \{i : \pi_i > \pi_{i+1} \text{ for } 0 \leq i < n\}$$

where $\pi_0 = -\pi_2$.

Example

$$\pi = 5\bar{7}2\bar{3}\bar{8}\bar{1}46 \in D_8$$

To find descent, we add a 7 in front, and calculate like “normal”.

$$75\bar{7}2\bar{3}\bar{8}\bar{1}46$$

$$012345678$$

$$\text{des}_D(\pi) = \emptyset$$

Descents - Type D

For $\pi = \pi_1 \dots \pi_n \in D_n$ let $\text{des}_D(\pi)$ denote the descent set of π .

$$\text{des}_D(\pi) = \{i : \pi_i > \pi_{i+1} \text{ for } 0 \leq i < n\}$$

where $\pi_0 = -\pi_2$.

Example

$$\pi = 5\bar{7}2\bar{3}\bar{8}\bar{1}46 \in D_8$$

$$75\bar{7}2\bar{3}\bar{8}\bar{1}46$$

$$012345678$$

$$\text{des}_D(\pi) = \{0\}$$

Descents - Type D

For $\pi = \pi_1 \dots \pi_n \in D_n$ let $\text{des}_D(\pi)$ denote the descent set of π .

$$\text{des}_D(\pi) = \{i : \pi_i > \pi_{i+1} \text{ for } 0 \leq i < n\}$$

where $\pi_0 = -\pi_2$.

Example

$$\pi = 5\bar{7}2\bar{3}\bar{8}\bar{1}46 \in D_8$$

75 $\bar{7}$ 2 $\bar{3}$ $\bar{8}$ $\bar{1}$ 46

012345678

$$\text{des}_D(\pi) = \{0, 1\}$$

Descents - Type D

For $\pi = \pi_1 \dots \pi_n \in D_n$ let $\text{des}_D(\pi)$ denote the descent set of π .

$$\text{des}_D(\pi) = \{i : \pi_i > \pi_{i+1} \text{ for } 0 \leq i < n\}$$

where $\pi_0 = -\pi_2$.

Example

$$\pi = 5\bar{7}2\bar{3}\bar{8}\bar{1}46 \in D_8$$

$$75\bar{7}2\bar{3}\bar{8}\bar{1}46$$

$$012\textcolor{red}{34}5678$$

$$\text{des}_D(\pi) = \{0, 1, 3\}$$

Descents - Type D

For $\pi = \pi_1 \dots \pi_n \in D_n$ let $\text{des}_D(\pi)$ denote the descent set of π .

$$\text{des}_D(\pi) = \{i : \pi_i > \pi_{i+1} \text{ for } 0 \leq i < n\}$$

where $\pi_0 = -\pi_2$.

Example

$$\pi = 5\bar{7}2\bar{3}\bar{8}\bar{1}46 \in D_8$$

$$75\bar{7}2\bar{3}\bar{8}\bar{1}46$$

$$0123\textcolor{red}{4}5678$$

$$\text{des}_D(\pi) = \{0, 1, 3, 4\}$$

Descents - Type D

For $\pi = \pi_1 \dots \pi_n \in D_n$ let $\text{des}_D(\pi)$ denote the descent set of π .

$$\text{des}_D(\pi) = \{i : \pi_i > \pi_{i+1} \text{ for } 0 \leq i < n\}$$

where $\pi_0 = -\pi_2$.

Example

$$\pi = 5\bar{7}2\bar{3}\bar{8}\bar{1}46 \in D_8$$

$$\text{des}_D(\pi) = \{0, 1, 3, 4\}$$

Main Theorem

Theorem (Bergeron, D., Machacek 2020BP)


The order complex $\Delta(P_n)$ is partitionable. Moreover,

$$h_i(\Delta(P_n)) = |\{\pi \in D_n : |\text{des}_B(\pi)| = i\}|$$

for each $0 \leq i \leq n$.

Example


$$h(\Delta(P_2)) = (1, 2, 1)$$



Permutations and maximal chains

How do we associate permutations and maximal chains in our poset?

- Change π_i to 0 inductively.



Permutations and chains

How do we associate permutations and chains in our poset?

- Order 0s first
- Add increasing sequences inductively.

57238146

Permutations and chains

How do we associate permutations and chains in our poset?

- Order 0s first
- Add increasing sequences inductively.

$$57|238|146 \xrightarrow{\min}$$

Permutations and chains

How do we associate permutations and chains in our poset?

- Order 0s first
- Add increasing sequences inductively.

$$57|238|146 \xrightarrow{\min} (+, +, +, +, 0, +, 0, +)$$

Permutations and chains

How do we associate permutations and chains in our poset?

- Order 0s first
- Add increasing sequences inductively.

$$57|238|146 \xrightarrow{\min} \begin{array}{c} (+, +, +, +, 0, +, 0, +) \\ | \\ (+, \textcolor{red}{0}, \textcolor{red}{0}, +, 0, +, 0, \textcolor{red}{0}) \end{array}$$

Permutations and chains

How do we associate permutations and chains in our poset?

- Order 0s first
- Add increasing sequences inductively.

$$57|238|\textcolor{red}{146} \xrightarrow{\min} \begin{array}{c} (+, +, +, +, 0, +, 0, +) \\ | \\ (\textcolor{red}{+}, 0, 0, \textcolor{red}{+}, 0, \textcolor{red}{+}, 0, 0) \end{array}$$

Permutations and chains

How do we associate permutations and chains in our poset?

- Order 0s first
- Add increasing sequences inductively.

$$\begin{array}{c} (+, +, +, +, 0, +, 0, +) \\ | \\ 57238146 \xrightarrow{\min} (+, 0, 0, +, 0, +, 0, 0) \end{array}$$

$$\text{des}_A(\pi) = \{2, 5\}$$

Permutations and chains

How do we associate permutations and chains in our poset?

- Order 0s first
- Add increasing sequences inductively.

$$\begin{array}{c} (+, +, +, +, 0, +, 0, +) \\ | \\ 57238146 \xrightarrow{\min} (+, 0, 0, +, 0, +, 0, 0) \end{array} \rightarrow$$

$$\text{des}_A(\pi) = \{2, 5\}$$

Permutations and chains

How do we associate permutations and chains in our poset?

- Order 0s first
- Add increasing sequences inductively.

$$\begin{array}{c} (+, +, +, +, \mathbf{0}, +, \mathbf{0}, +) \\ | \\ 57238146 \xrightarrow{\min} (+, 0, 0, +, 0, +, 0, 0) \rightarrow 57 \end{array}$$

$$\text{des}_A(\pi) = \{2, 5\}$$

Permutations and chains

How do we associate permutations and chains in our poset?

- Order 0s first
- Add increasing sequences inductively.

$$\begin{array}{c} (+, +, +, +, 0, +, 0, +) \\ | \\ 57238146 \xrightarrow{\min} (+, \textcolor{red}{0}, \textcolor{red}{0}, +, 0, +, 0, \textcolor{red}{0}) \end{array} \rightarrow 57238$$

$$\text{des}_A(\pi) = \{2, 5\}$$

Permutations and chains

How do we associate permutations and chains in our poset?

- Order 0s first
- Add increasing sequences inductively.

$$\begin{array}{c} (+, +, +, +, 0, +, 0, +) \\ | \\ 57238146 \xrightarrow{\min} (+, 0, 0, +, 0, +, 0, 0) \rightarrow 57238146 \end{array}$$

$$\text{des}_A(\pi) = \{2, 5\}$$

Permutations and chains

How do we associate permutations and chains in our poset?

- Order 0s first
- Add increasing sequences inductively.

$$\begin{array}{c} (+, +, +, +, 0, +, 0, +) \\ | \\ 57238146 \xrightarrow{\min} (+, 0, 0, +, 0, +, 0, 0) \rightarrow 57238146 \end{array}$$

$$\text{des}_A(\pi) = \{2, 5\}$$

Negatives?

But how do we handle the negatives?!

$$5\bar{7}2\bar{3}\bar{8}\bar{1}46 \leftrightarrow ?$$

Negatives?

But how do we handle the negatives?!

$$\begin{array}{c} \bar{5} \bar{7} 2 \bar{3} \bar{8} \bar{1} 4 6 \\ \leftrightarrow \\ ? \end{array}$$

(57238146, {1, 3, 7, 8})

Sign Variations

Sign vector $\omega \in \mathcal{V}_n$.

$\text{var}(\omega)$ = number of times ω changes sign

$i \in [n]$ is a *sign flip* of ω if there exists a j such that $\omega_{i-j}\omega_i < 0$ while $\omega_{i-k}\omega_i = 0$ for all $1 \leq k < j$.

Example

$$\omega = (+, +, -, -, -, -, +, -) \Rightarrow \text{var}(\omega) = 3$$

$$(+, +, -, -, -, 0, +, -) \leftrightarrow \{3, 7, 8\}$$

1 2 3 4 5 6 7 8

Cyclic Sign Variations


Sign vector $\omega \in \mathcal{V}_n$.

$cvar(\omega)$ = number of times ω changes sign, cyclically

$i \in [n]$ is a *cyclic sign flip* of ω if there exists a j such that $\omega_{i-j}\omega_i < 0$ while $\omega_{i-k}\omega_i = 0$ for all $1 \leq k < j$ where $\omega_i = \omega_{i+n}$.

Example


$$\omega = (+, +, -, -, -, -, +, -) \Rightarrow cvar(\omega) = 4$$



Even signed permutations and maximal chains

How about even signed permutations and maximal chains?

- Just as before with a final step of adding negatives.



Even signed permutations and chains

How do we associate even signed permutations and chains in our poset?

- Just as before, but with negatives.

5 $\bar{7}$ 2 $\bar{3}$ 8 $\bar{1}$ 46

(57238146, {1, 3, 7, 8})

Even signed permutations and chains

How do we associate even signed permutations and chains in our poset?

- Just as before, but with negatives.

$$5|\bar{7}2|\bar{3}|\bar{8}\bar{1}46 \xrightarrow{\min} \\ (57238146, \{1, 3, 7, 8\})$$

Even signed permutations and chains

How do we associate even signed permutations and chains in our poset?

- Just as before, but with negatives.

$$(+, +, -, -, -, -, +, -)$$

$$\begin{array}{c} 5|\bar{7}2|\bar{3}|\bar{8}\bar{1}46 \xrightarrow{\min} \\ (57238146, \{1, 3, 7, 8\}) \end{array}$$

Even signed permutations and chains

How do we associate even signed permutations and chains in our poset?

- Just as before, but with negatives.

$$(+, +, -, -, -, -, +, -)$$

$$\begin{array}{c} 5| \bar{7}2|\bar{3}|\bar{8}\bar{1}46 \xrightarrow{\min} (+, +, -, -, \mathbf{0}, -, +, -) \\ (\textcolor{red}{57238146}, \{1, 3, 7, 8\}) \end{array}$$

Even signed permutations and chains

How do we associate even signed permutations and chains in our poset?

- Just as before, but with negatives.

$$(+, +, -, -, -, -, +, -)$$

$$\begin{array}{c} 5|\bar{7}\bar{2}|\bar{3}|\bar{8}\bar{1}46 \xrightarrow{\min} (+, +, -, -, 0, -, +, -) \\ (\textcolor{red}{57238146}, \{1, 3, 7, 8\}) \qquad \qquad (+, \textcolor{red}{0}, -, -, 0, -, \textcolor{red}{0}, -) \end{array}$$

Even signed permutations and chains

How do we associate even signed permutations and chains in our poset?

- Just as before, but with negatives.

$$(+, +, -, -, -, -, +, -)$$

$$\begin{array}{ccc} 5|\bar{7}2|\bar{\textcolor{red}{3}}|\bar{8}\bar{1}46 & \xrightarrow{\min} & (+, +, -, -, 0, -, +, -) \\ (\textcolor{red}{57238146}, \{1, 3, 7, 8\}) & & | \\ & & (+, 0, -, -, 0, -, 0, -) \\ & & | \\ & & (+, 0, \textcolor{red}{0}, -, 0, -, 0, -) \end{array}$$

Even signed permutations and chains

How do we associate even signed permutations and chains in our poset?

- Just as before, but with negatives.

$$(+, +, -, -, -, -, +, -)$$

$$\begin{array}{c} 5|\bar{7}2|\bar{3}|\bar{\bar{8}}\bar{1}46 \xrightarrow{\min} (+, +, -, -, 0, -, +, -) \\ (\textcolor{red}{57238146}, \{1, 3, 7, 8\}) \qquad \qquad \qquad | \\ \qquad \qquad \qquad (+, 0, -, -, 0, -, 0, -) \\ \qquad \qquad \qquad | \\ \qquad \qquad \qquad (\textcolor{red}{+, 0, 0, -, 0, -, 0, -}) \end{array}$$

Even signed permutations and chains

How do we associate even signed permutations and chains in our poset?

- Just as before, but with negatives.

$$(+, +, -, -, -, -, +, -)$$

$$\begin{array}{c} (+, +, -, -, 0, -, +, -) \\ \downarrow \\ (+, 0, -, -, 0, -, 0, -) \\ \downarrow \\ (+, 0, 0, -, 0, -, 0, -) \\ \text{des}_D(\pi) = \{0, 1, 3, 4\} \end{array}$$

$5|\bar{7}2|\bar{3}|\bar{8}\bar{1}46 \xrightarrow{\min}$

$(57238146, \{1, 3, 7, 8\})$

Even signed permutations and chains

How do we associate even signed permutations and chains in our poset?

- Just as before, but with negatives.

$$(+, +, -, -, -, -, +, -)$$

$$\begin{array}{c} (+, +, -, -, 0, -, +, -) \\ | \\ (+, 0, -, -, 0, -, 0, -) \\ | \\ (+, 0, 0, -, 0, -, 0, -) \\ \text{des}_D(\pi) = \{0, 1, 3, 4\} \\ \text{des}_B(\pi) = \{1, 3, 4\} \end{array}$$

5| $\bar{7}2|\bar{3}| $\bar{8}146 $\xrightarrow{\min}$ (57238146, {1, 3, 7, 8})$$

Even signed permutations and chains

How do we associate even signed permutations and chains in our poset?

- Just as before, but with negatives.

$$\begin{array}{c} (+, +, -, -, 0, -, +, -) \\ | \\ (+, 0, -, -, 0, -, 0, -) \rightarrow \\ | \\ (+, 0, 0, -, 0, -, 0, -) \end{array}$$

$\overline{5} \overline{7} 2 \overline{3} \overline{8} \overline{1} 4 6 \xrightarrow{\min}$

Even signed permutations and chains

How do we associate even signed permutations and chains in our poset?

- Just as before, but with negatives.

$$\begin{array}{c} (+, +, -, -, 0, -, +, -) \\ | \\ (+, 0, -, -, 0, -, 0, -) \rightarrow \{1, 3, 7, 8\} \\ | \\ (+, 0, 0, -, 0, -, 0, -) \end{array}$$

5\bar{7}2\bar{3}\bar{8}\bar{1}46 \xrightarrow{\min}

Even signed permutations and chains

How do we associate even signed permutations and chains in our poset?

- Just as before, but with negatives.

$$\begin{array}{c} (+, +, -, -, \textcolor{red}{0}, -, +, -) \\ | \\ \{1, 3, 7, 8\} \\ 5\bar{7}2\bar{3}\bar{8}\bar{1}46 \xrightarrow{\min} (+, 0, -, -, 0, -, 0, -) \rightarrow \\ | \\ (+, 0, 0, -, 0, -, 0, -) \end{array}$$

5

Even signed permutations and chains

How do we associate even signed permutations and chains in our poset?

- Just as before, but with negatives.

$$\begin{array}{c} (+, +, -, -, 0, -, +, -) \\ | \\ \{1, 3, 7, 8\} \\ 5\bar{7}2\bar{3}\bar{8}\bar{1}46 \xrightarrow{\min} (+, \textcolor{red}{0}, -, -, 0, -, \textcolor{red}{0}, -) \rightarrow \\ | \\ (+, 0, 0, -, 0, -, 0, -) \end{array}$$

5\bar{7}2

Even signed permutations and chains

How do we associate even signed permutations and chains in our poset?

- Just as before, but with negatives.

$$\begin{array}{c} (+, +, -, -, 0, -, +, -) \\ | \\ \{1, 3, 7, 8\} \\ 5\bar{7}2\bar{3}\bar{8}\bar{1}46 \xrightarrow{\min} (+, 0, -, -, 0, -, 0, -) \rightarrow \\ | \\ (+, 0, \textcolor{red}{0}, -, 0, -, 0, -) \\ 5\bar{7}2\bar{3} \end{array}$$

Even signed permutations and chains

How do we associate even signed permutations and chains in our poset?

- Just as before, but with negatives.

$$\begin{array}{c} (+, +, -, -, 0, -, +, -) \\ | \\ \{1, 3, 7, 8\} \\ 5\bar{7}2\bar{3}\bar{8}\bar{1}46 \xrightarrow{\min} (+, 0, -, -, 0, -, 0, -) \rightarrow \\ | \\ (+, 0, 0, -, 0, -, 0, -) \\ | \\ 5\bar{7}2\bar{3}\bar{8}\bar{1}46 \end{array}$$

Even signed permutations and chains

How do we associate even signed permutations and chains in our poset?

- Just as before, but with negatives.

$$\begin{array}{c} (+, +, -, -, 0, -, +, -) \\ | \\ (+, 0, -, -, 0, -, 0, -) \rightarrow \\ | \\ (+, 0, 0, -, 0, -, 0, -) \\ \hline 5\bar{7}2\bar{3}\bar{8}\bar{1}46 \xrightarrow{\min} (+, 0, -, -, 0, -, 0, -) \rightarrow 5\bar{7}2\bar{3}\bar{8}\bar{1}46 \end{array}$$

Main Theorem (Again)

Let D_n be the type D Coxeter group and let des_B denote the type B descent set of an element $\pi \in D_n$.

Theorem (Bergeron, D., Machacek 2020BP)

The order complex $\Delta(P_n)$ is partitionable. Moreover,

$$h_i(\Delta(P_n)) = |\{\pi \in D_n : |\text{des}_B(\pi)| = i\}|$$


for each $0 \leq i \leq n$.

Sign Variations and Descents

Example - $n = 2$

$$h_i(\Delta(P_2)) = |\{\pi \in D_2 : |\text{des}_B(\pi)| = i\}| \quad h(\Delta(P_2)) = (1, 2, 1)$$

D_n	des_B
12	\emptyset
$\bar{2}1$	{0}
21	{1}
$\bar{1}\bar{2}$	{0, 1}



Restriction of variations

- $\mathcal{PV}_{n,m} = \{\omega \in \mathcal{PV}_n : \text{var}(\omega) \leq m\}.$
- $P_{n,m} = (\mathcal{PV}_{n,m}, <).$
- $D_{n,m} = \{\pi \in D_n : \pi \text{ has at most } m \text{ negatives}\}.$

Theorem (Bergeron, D., Machacek 2020BP)

If $m \leq n - 1$ is even then the order complex $\Delta(P_{n,m})$ is partitionable. Moreover,

$$h_i(\Delta(P_{n,m})) = |\{\pi \in D_{n,m} : |\text{des}_B(\pi)| = i\}|$$

for each $0 \leq i \leq n$.

Sign Variations and Descents

Thank you!

$$\Delta(P_3)$$

